



## Congenital First Pharyngeal Arch Anomaly in a Holstein Friesian Calf

Bahador Shojaei<sup>a</sup>, Sina Delshad<sup>a</sup>, Mehdi Farajli Abbasi<sup>b</sup>, Maehdi Jaaferi<sup>c</sup>, Mohammad Farajli Abbasi<sup>d</sup>

<sup>a</sup> Department of Basic Sciences, Faculty of Veterinary Medicine, Shahid Bahonar University of Kerman, Iran.

<sup>b</sup> Doctor of Veterinary Medicine, Mehr Pet Clinic, Tabriz, Iran.

<sup>c</sup> Department of Clinical Sciences, School of Veterinary Medicine, Shahid Bahonar University of Kerman, Iran.

<sup>d</sup> Neuroscience Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Kerman, Iran

### ABSTRACT

The developmental error of the first pair of pharyngeal arch processes would result in a group of malformations related to the oral cavity. These malformations can be listed as agnathia, micrognathia, brachygnathia, syngnathia, cleft palate, campylognathia, parrot beak, and strophocephalia. Campylognathia is an anomaly in which the lower or upper jaw deviated from the midline. Clinical examination of a newborn female Holstein Friesian calf showed mandibular deviation, inferior and superior cheek teeth deformation, and some other facial malformations. No similar case was detected in the history of the herd. Precise consideration of malformed animals and the extent of involvement of different structures indicates the pattern of malformation, the time of teratogenic effect, and the motive embryonic structures. In the present case, the involvement of derivatives of both maxillary and mandibular processes shows the responsibility of the first pharyngeal arch for mentioned abnormalities. In humans, different types of abnormal development of the first pharyngeal arch have been reported and so, have been classified under the “first visceral arch syndrome”. But according to the infrequency of reports, this classification has not been done in domestic animals.

### Keywords

*Congenital cleft jaw, pharyngeal arch, calf, anomaly, campylognathia*

### Abbreviations

TRP: Temperature, Pulse, Respiratory rate  
BMP: Bone Morphogenetic Protein  
FGF: Fibroblast Growth Factor  
SHH: Sonic hedgehog  
WNT: Wingless-related integration site

Number of Figures: 2  
Number of Tables: 0  
Number of References: 21  
Number of Pages: 5

## Introduction

The mesenchyme of the first pair of pharyngeal arches gives rise to dorsal maxillary and ventral mandibular prominences. These facial structures converge and fuse in the midline enclosing an invagination of ectoderm, the stomodeum. The ventral mandibular prominences contribute to the formation of the lower jaw while the paired maxillary prominences procedure the upper jaw [1]. The developmental error of these structures, often due to the lack of migration of neural crest cells to the first pharyngeal arch, leads to a group of malformations related to the oral cavity [2]. The developmental error of these structures would result in a group of malformations related to the oral cavity. These malformations can be listed as agnathia, micrognathia, brachygnathia, Syngnathia, cleft palate, campylognathia [3,4], parrot beak, and strophocephalia [5]. Campylognathia is an anomaly in which the lower or upper jaw deviated from the midline [6]. It has been reported in geese [7], horses [8], sheep [4], and cows [9]. Lack of maternal protein and vitamins, infectious diseases, radiation, folic acid deficiency, teratogens, and endocrine disorders at the beginning of organogenesis lead to the formation of these malformations [10,11]. In humans, this anomaly has been reported as one of several malformations of the "First Arch Syndrome". But, in animals, no classification has been made on the pharyngeal arch abnormalities. In the present work, we aim to describe a Holstein Friesian calf with severe inferior campylognathia associated with the cleft palate and oligodontia from the Kerman province for the first time.

## Case Presentation


A seven-day-old newborn female Holstein Friesian calf was referred to the veterinary hospital, Shahid Bahonar University of Kerman. On the clinical examination, mandibular deviation, respective upper and lower cheek teeth deformation, and malformation of the buccal and masseter regions on the left side of the face was seen (Figure 1). Regarding the history of the herd, it was the fifth delivery of the mother and the previous deliveries were normal, this was the first case in the herd, also no sign of poisoning was found and no medicine was used during pregnancy. TRP (Temperature, Pulse, Respiratory rate), and other vital signs have been normal. The calf had fed her mother milk. There was no symptom of respiration pneumonia, but milk was fed flow out from both nostrils during feeding. According to the severity of malformations, the animal was sacrificed and dissected. The necropsy was revealed a severe mandibular deviation to the left side

of the face (Figure 2A). The ramus on the left side was short without any remnant of the condylar process (Figure 2B). The ipsilateral masseter muscle has a very poor configuration. The origin of the masseter on the facial bones was distorted and its insertion was seen as sparse fascicles on the abnormal mandibular ramus (Figure 2C). According to the shortening of the ramus, the caudal portion of the medial pterygoids was exposed. Both mandibular and maxillary oligodontia and Crooked teeth were seen in the right and left jaws. The rostral part of the hard palate was intact but a longitudinal cleft extended from the dental pad caudally to the pharyngeal region to involve both hard and soft palates (secondary cleft palate). The right external ear was intact. The left auricle has been formed but no external auditory meatus was detected. The left mandibular ramus was narrow and deformed. No other abnormalities were observed in the dissection of the brain and other organs. Also, the absence of ataxia, neurological symptoms such as turning around, head pressing, or any clinical neurological symptoms indicated the absence of this type of brain involvement and confirmed the anatomical evidence.



**Figure 1.**  
A) Left view of the face of affected animal. B) Malformed buccal region results extension of the oral commissure nearby the external ear.

First Pharyngeal Arch Anomaly



**Figure 2.**  
A) Mandibular campylognathia. B) Secondary cleft palate (arrows). C) Exposition of the medial pterygoideus according to the malformation in the mandible. The oligodontia is also seen in the mandibular (arrowed A) and maxillary (arrowed B).

First Pharyngeal Arch Anomaly



## Results & Discussion

According to the interaction of several organ systems, the morphogenesis of the face seems very complicated [12]. It is known that the fetal organs such as the facial processes, the pharyngeal arches, and the developing brain, would deeply relate to the formation of the face in early embryonic life [12,13]. So, in the evaluation of the facial anomalies, attention should be paid to the derivatives of each embryonic precursor. Precise consideration of the malformed animals and the extent of involvement of different facial structures, revealed the pattern of malformation, the time of the effect of teratogen, and the motive embryonic structures [14]. In the present case, all of the detected disorders incriminate the first pharyngeal arch. Secondary cleft palate deviated mandible and its rudimentary ramus, malformed masticatory muscles, and oligodontia support the involvement of the maxillary and mandibular processes of this arch. Clefts of the face are developmental disorders due to failure of closure in facial processes such as the frontonasal, maxillary, and mandibular processes. They can be asymmetric unilateral or median symmetrical clefts. In the ruminants, the most reported type of these disorders is the secondary cleft palate, which is shaped due to failure of the growth of the maxillary processes of the first pharyngeal arches, but cleft palate accompanying with the campylognathia reported as a rare anomaly in the herbivores [8]. In the present case involvement of derivatives of both maxillary and mandibular processes shows the responsibility of the first pharyngeal arch for mentioned abnormalities. In humans, different types of abnormal development of the first pharyngeal arch have been reported and so, have been classified under the "first visceral arch syndrome" [15]. But according to the infrequency of reports, this classification has not been done in domestic animals.

Since the growth of teeth and the development of the craniofacial structures are affected by similar biological factors such as BMP, FGF, SHH, and WNT [16,17], therefore in many abnormal cases, malformation of these structures are accompanied by each other, as we saw in this case.

The critical period of fusion of facial fissure and palate took place at days 34 and 56 of gestation, respectively [18]. Although a genetic basis has been determined for the cleft palate (multifactorial or autosomal inheritance in the Charolais breed) and mandibular deformations (brachygnathia inferior) (polygenic inheritance in the Simmental breed) [8,19,20] as we did not find any similar case in the herd history and previous parturitions, also there was evidence of no poisoning and no change in diet, it may be stated that the present case has been exposed to environ-

mental teratogens such as poisonous plants and viruses (e.g. Bluetongue virus) in the first two months of embryogenesis [21].

## Ethics statement

This study was approved by the research ethics committee of the Shahid Bahonar University of Kerman, Iran (IR.UK.VETMED.REC). Animal care and all procedures were accomplished according to the instructions of care and use of laboratory animals.

## Authors' Contributions

BS and SD performed review literature and manuscript writing. MFA and MFA performed the clinical examination and necropsy MJ examined the animal and performed review literature.

## Competing Interests

The authors declare that there is no conflict of interest.

## References

- McGeady TA, Quinn PJ, Fitzpatrick ES, Ryan MT, Kilroy D, Lonergan P. Veterinary embryology. 2nd ed. John Wiley & Sons; 2017. 274–288.
- The Developing Human: Clinically Oriented Embryology, 9e - Keith L. Moore - Google Books [Internet]. 2019 [cited 2021 Jul 11]. p. 143–79. Available from: [https://books.google.com/books?hl=en&lr=&id=OTaBDwAAQBAJ&oi=fnd&pg=PP1&ots=GoyHv9ri7z&sig=29Ch-4A6\\_MpoH2PtxDCKZt-TH3QU#v=onepage&q&f=false](https://books.google.com/books?hl=en&lr=&id=OTaBDwAAQBAJ&oi=fnd&pg=PP1&ots=GoyHv9ri7z&sig=29Ch-4A6_MpoH2PtxDCKZt-TH3QU#v=onepage&q&f=false)
- Flueck WT, Smith-Flueck JAM. Diseases of red deer introduced to Patagonia and implications for native ungulates. *Anim Prod Sci*. 2012; 52(8):766–73.
- Shojaei B, Derakhshanfar A, Oloumi MM, Hashemnia S. Di-prosopus, spina bifida and kyphoscoliosis in a lamb-a case report. *Vet Arh*. 2006; 76(5): 461–9.
- Lejong M, Vanmuylder N, Louryan S. External and computed tomography analysis of a strophocephalic lamb. *Morphologie*. 2019; 103(341):122–5.
- Shojaei B, Asadi A. Dicephalus, inferior brachygnathia, cleft palate and campylognathia in a calf. *Iran J Vet Res*. 2008; 4:117–22.
- Lingnau A, Kaleta EF. Malformations of the upper beak (campylognathia superior) and adjacent bones within the splanchnocranium of domestic geese: an evaluation of cases that have been described during four centuries. *Vet Hist*. 2018; 19(2):163–76.

- Flueck WT, Smith-Flueck JAM. Huemul heresies: beliefs in search of supporting data. 2. Biological and ecological considerations. *Anim Prod Sci*. 2012; 52(8): 694–706.
- Leipold HW, Ojo SA, Huston K. Genetic Defects of the Skeletal System in Cattle. In: 1st World Congress on Genetics Applied to Livestock Production: Symposia. 1974; 35–40.
- Hobbenaghi R, Dalir-Naghadeh B, Nazarizadeh A. Congenital multi-organ malformations in a Holstein calf. *Vet Res Forum*. 2015; 6(3): 257–260.
- Oryan A, Shirian S, Samadian M.R. Congenital craniofacial and skeletal defects with arthrogryposis in two newborn male Holstein Friesian calves. *Comp Clin Pathol*. 2011; 20: 43–46.
- Pruzansky S. Anomalies of face and brain. *Birth Defects Orig Artic Ser*. 1975; 11(7):183–204.
- DeMyer W. Median facial malformations and their implications for brain malformations. *Birth Defects Orig Artic Ser*. 1975; 11(7): 155–81.
- Moritomo Y, Tsuda T, MIYAMOTO H. Craniofacial skeletal abnormalities in anomalous calves with clefts of the face. *J Vet Med Sci*. 1999; 61(10): 1147–52.
- McKenzie J. The first arch syndrome. *Arch Dis Child*. 1958; 33(171):477.
- Seppala M, Fraser GJ, Birjandi AA, Xavier GM, Cobourne MT. Sonic hedgehog signaling and development of the dentition [Internet]. Vol. 5, *Journal of Developmental Biology: Multidisciplinary Digital Publishing Institute*; 2017 [cited 2021 Jul 11]; 6. Available from: <https://www.mdpi.com/2221-3759/5/2/6/htm>
- Cobourne MT, Sharpe PT. Diseases of the tooth: The genetic and molecular basis of inherited anomalies affecting the dentition [Internet]. Vol. 2, *Wiley Interdisciplinary Reviews: Developmental Biology*. John Wiley & Sons, Ltd; 2013 [cited 2021 Jul 11]; 183–212. Available from: <https://wires-onlinelibrary-wiley-com.ezp4.semantak.com/doi/full/10.1002/wdev.66>
- Dixon MJ. Treacher Collins syndrome. *J Med Genet*. 1995; 32(10):806.
- Evans HE, WO S. Prenatal development of domestic and laboratory mammals: growth curves, external features and selected references. *Anat Histol Embryol*. 1973; 2:11–45.
- Rousseaux CG. Congenital defects as a cause of perinatal mortality of beef calves. *Vet Clin North Am Food Anim Pract*. 1994; 10(1): 35–51.
- Line BS. Congenital and Inherited Anomalies in Animals. 2020 [cited 2021 Jul 10];1–8. Available from: <https://www.msdevetmanual.com/generalized-conditions/congenital-and-inherited-anomalies/congenital-and-inherited-anomalies-in-animals>

## COPYRIGHTS

©2022 The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.



## How to cite this article

Shojaei B, Delshad S, Farajili Abbasi M, Jafari M, Farajili Abbasi M. Congenital First Pharyngeal Arch Anomaly in a Holstein Friesian Calf. *Iran J Vet Sci Technol*. 2022; 14(1):45–49.  
DOI: <https://doi.org/10.22067/ijvst.2022.73253.1094>  
URL: [https://ijvst.um.ac.ir/article\\_42069.html](https://ijvst.um.ac.ir/article_42069.html)